

UT Southwestern Department of Radiology

Protocol Name: CT FAI - Unilateral
Epic Button: CT FAI - Unilateral
Indications: CT evaluation for unilateral femoroacetabular impingement and/or hip dysplasia, femoral torsion

Orderable Name: CT LOWER EXTREMITY LEFT WO IV CONTRAST
CT LOWER EXTREMITY RIGHT WO IV CONTRAST

Adult Only

CTDIvol < 50 mGy

Acquisitions: 2

Active Protocol

Oral Contrast: None	IV Contrast: None	Other Contrast: None	Airway
			Other Notes Patient supine, tape toes together Scan through pelvis/hips, then scan knee from side of complaint (cover femoral condyles.) See HipMap Scan Protocol link for details. Use both RIGHT and LEFT orderables. If the patient only has 1 hip, use only the appropriate RIGHT or LEFT orderable.

Last Change: 11/4/2025

Last Review: 12/4/2025

Links: [General Statements](#) [FAI HipMap Scan Guide](#)

Special Instructions		Do not repeat CT scan, recon from 1st acquisition	Do not repeat CT scan, recon from 1st acquisition	Do not repeat CT scan, recon from 1st acquisition
Acq # / Series Name	1 Noncontrast	N/A Noncontrast	N/A Noncontrast	N/A Noncontrast
Phase Timing			N/A	
Acquisition Protocol		Recon Only	Recon Only	Recon Only
Coverage	Iliac crest to below lesser trochanter	Iliac crest to below lesser trochanter	Acetabular roof to below lesser trochanter	Acetabular roof to below lesser trochanter
FOV	Pelvis	Pelvis	Right Hip	Right Hip
Algorithm	Soft Tissue	Bone	Soft Tissue	Bone
Axial Recons	3 mm, Volume	3 mm, Volume	3 mm	3 mm
Other Planar Recons	3 mm coronal and sagittal	3 mm coronal and sagittal	3 mm coronal and sagittal	3 mm coronal and sagittal
MIP Recons				
†DECT Philips	Gout maps (cor/sag), BM edema, SBI			
†DECT Siemens	Gout maps (cor/sag), BM edema, low/high kVp, mono E 100, mono E 120			
†PC-CT Siemens				

UT Southwestern Department of Radiology

Protocol Name: CT FAI - Unilateral
Epic Button: CT FAI - Unilateral
Indications: CT evaluation for unilateral femoroacetabular impingement and/or hip dysplasia, femoral torsion

Orderable Name: CT LOWER EXTREMITY LEFT WO IV CONTRAST
CT LOWER EXTREMITY RIGHT WO IV CONTRAST

Adult Only
CTDIvol < 50 mGy

Acquisitions: 2

Active Protocol

Oral Contrast: None	IV Contrast: None	Other Contrast: None	Airway
			Other Notes Patient supine, tape toes together Scan through pelvis/hips, then scan knee from side of complaint (cover femoral condyles.) See HipMap Scan Protocol link for details. Use both RIGHT and LEFT orderables. If the patient only has 1 hip, use only the appropriate RIGHT or LEFT orderable.

Last Change: 11/4/2025

Last Review: 12/4/2025

Links: [General Statements](#) [FAI HipMap Scan Guide](#)

Special Instructions	Do not repeat CT scan, recon from 1st acquisition	Do not repeat CT scan, recon from 1st acquisition	
Acq # / Series Name	N/A Noncontrast	N/A Noncontrast	2 Noncontrast
Phase Timing			
Acquisition Protocol	Recon Only	Recon Only	
Coverage	Acetabular roof to below lesser trochanter	Acetabular roof to below lesser trochanter	Inferior patella to top of tibial plateau
FOV	Left Hip	Left Hip	Affected Knee
Algorithm	Soft Tissue	Bone	Soft Tissue
Axial Recons	3 mm	3 mm	3 mm, Volume
Other Planar Recons	3 mm coronal and sagittal	3 mm coronal and sagittal	
MIP Recons			
†DECT Philips			
†DECT Siemens			
†PC-CT Siemens			

† When dual energy (DE) or photon counting (PC) CT is used

